

كيسثوالفا

(الفاكيثواينالوك)

(Alpha Keto Analogue)

Composition:

Each film coated tablet contains:

α -Ketoanalogue to isoleucine (Calcium salt)	67mg
α -ketoanalogue to leucine (Calcium-salt)	101mg
lpha-ketoanalogue to phenylalanine (Calcium-salt)	68 mg
lpha-ketoanalogue to valine (Calcium-salt)	86 mg
lpha-ketoanalogue to methionine (Calcium-salt)	59 mg
L-Lysine-acetate	105 mg
L-Threonine	53 mg
L-Tryptophan	23 mg
L-Histidine	38 mg
L-Tyrosine	30 mg
Total Nitrogen content per tab 36mg, calcium per ta	b 0.05gm
(Genome Specs)	

Indications:

Prevention and therapy of damage due to faulty or deficient protein metabolism in chronic renal insufficient in connection with limited protein in food of 40 gram per day (for adults) and less; i.e generally in patients with a glomerular filtration rate (GFR) below 25 ml/min.

Contra-Indications:

Hypercalcemia, disturbed amino acid metabolism In case of hereditary phenylketonuria it has taken Into account that this product contain phenylalanine

Note:

No experience has been made so far with the application in pregnancy and paediatrics.

Side-effects:

Hpercalcemia may develop.

Note:

Monitor regularly the serum calcium level.

Interactions with other drugs:

The simultaneous administration of medicaments containing calcium (e.g. acetolyte) may lead to pathological increase of the serum calcium level or intensifi-cation. As the uremic symptoms improve under ketoalfa a possible administration of aluminium hydroxide should be reduced. Pay attention to a reduction of serum phosphate. In order not to interfere with absorption, do not take drugs togather with ketoalfa that form sparingly soluble compounds with calcium (e.g.

tetracyclines, quinolones like Ciprofloxacin and Norfloxacin, iron, Fluoride and Estramustin-containing drugs). Between the intake of Ketoalfa tablets and Drugs from the mentioned categories a period of at least 2 hours should pass. If the blood calcium concentration raises by the intake of Ketoalfa tablets the susceptibility towards certain drugs enhancing the heart strength (heart/cardiac-active glycosides) will increase and the risk of heart rhythum disturbance will increase.

Application and Dosage:

For oral use.

If not otherwise prescribed, take 4 to 8 tablets three times a day during meals. Swallow whole. This dosage applies to adults (70 kg/BW).

Duration of Application:

Ketoalfa tablets are given as long as the glomerular filtration rate is below25ml/minute. Simultaneously food should contain 40 g/day protein or less (for adults).

Precautions:

Keep out of the reach of children! Do not use Ketoalfa tablets after expiry date!

Presentation:

Pack containing 10x10's Tablets

Storage Condition:

Store below 30°C.

Protect from heat, moisture & sunlight.

خوراک: دواڈ اکٹر کی ہدایت کے مطابق استعال کریں۔ احتیاط: دواکودھوپ،نمی اور گرمی سے بچائیں۔ تمام ادویات بچوں کی پہنچ سے دورر کھیں۔ دواکو معاڈ گری سینٹی گریڈسے کم درجہ حررات پر کھیں۔



